arXiv:0809.0933v1 [hep-lat] 4 Sep 2008

Modern Physics Letters A
© World Scientific Publishing Company

PROPERTIES OF SCALAR-QUARK SYSTEMS IN SU(3)$_c$ LATTICE QCD

HIDEAKI IIDA and TORU T. TAKAHASHI
Yukawa Institute for Theoretical Physics (YITP), Kyoto University
Kitashirakawaoiwake, Sakyo, Kyoto 606-8502, Japan
iida@yukawa.kyoto-u.ac.jp

HIDEO SUGANUMA
Department of Physics, Kyoto University
Kitashirakawaoiwake, Sakyo, Kyoto 606-8502, Japan

Received 17 February 2008
Revised (Day Month Year)

We perform the first study for the bound states of colored scalar particles ϕ ("scalar quarks") in terms of mass generation with quenched SU(3)$_c$ lattice QCD. We investigate the bound states of ϕ, $\phi^\dagger\phi$ and $\phi\phi\phi$ ("scalar-quark hadrons"), as well as the bound states of ϕ and quarks ψ, i.e., $\phi^\dagger\psi$, $\psi\psi\phi$ and $\phi\phi\psi$ ("chimera hadrons"). All these new-type hadrons including ϕ have a large mass of several GeV due to large quantum corrections by gluons, even for zero bare scalar-quark mass $m_\phi = 0$ at $a^{-1} \sim 1$GeV. We find a similar m_ψ-dependence between $\phi^\dagger\psi$ and $\phi\phi\psi$, which indicates their similar structure due to the large mass of ϕ. From this study, we conjecture that all colored particles generally acquire a large effective mass due to dressed gluons.

Keywords: Dynamical mass generation; Lattice QCD; Scalar-quarks; Diquarks.

PACS Nos.: 12.38.Gc, 12.38.Mh, 14.40.Gx, 25.75.Nq

1. Introduction

The origin of mass is one of the fundamental and fascinating subjects in physics. About 99% of mass of matter in the world originates from the strong interaction, which provides the large constituent quark mass $M_\psi = (300-400)$MeV. Such a dynamical fermion-mass generation in the strong interaction can be interpreted as spontaneous chiral-symmetry breaking (χSB).[1,2]

In the strong interaction, however, there is other type of dynamical mass generation than χSB. For instance, gluons, which are massless in perturbation QCD, seem to have a large effective mass as $(0.5-1.0)$GeV,[3,4] due to non-perturbative effects. Actually, glueballs, which are ideally composed only by gluons, have a large mass, e.g., about 1.5GeV.[5,6] The same holds for charm quarks. Whereas the current mass of charm quarks is about 1.2GeV at the renormalization point $\mu = 1$GeV,

Table 1. The lattice QCD setup for scalar-quark hadrons and chimera hadrons.

β	lattice size	lattice spacing a	bare scalar-quark mass m_ϕ	bare quark mass m_ψ
5.7	$16^3 \times 32$	$(1.1\mathrm{GeV})^{-1} \simeq 0.2\mathrm{fm}$	0.00, 0.11, 0.22, 0.33GeV	0.09, 0.14, 0.19GeV

the constituent charm-quark mass in the quark model is set to be about 1.6GeV. The about 400MeV difference between the current and the constituent charm-quark masses could be explained by dynamical mass generation without χSB, since there is no chiral symmetry for such heavy quarks. These examples imply mass generation without χSB in the strong interaction. We therefore conjecture that large dynamical mass generation generally occurs even without χSB in the strong-interaction world, i.e., *all colored particles have a large effective mass generated by dressed gluon effects.*[9,10] In this study, we investigate the system of colored scalar particles, which do not have chiral symmetry.

2. Scalar-quark Hadrons and Chimera Hadrons in Lattice QCD

We consider light 3c-colored "scalar-quarks" ϕ. The light scalar-quarks can be also regarded as idealized point-like "diquarks" at the scale of $a^{-1} \sim 1$GeV. We investigate "scalar-quark mesons" $\phi^\dagger\phi$ and "scalar-quark baryons" $\phi\phi\phi$ as the bound states of scalar quarks ϕ. We also investigate the bound states of scalar-quarks ϕ and quarks ψ, i.e., $\phi^\dagger\psi$, $\psi\psi\phi$ and $\phi\phi\psi$, which we name "chimera hadrons".[9,10]

To include scalar-quarks ϕ together with quarks ψ and gluons in QCD, we adopt the generalized QCD Lagrangian density,[9,10]

$$\mathcal{L} = -\frac{1}{4} G^a_{\mu\nu} G^{a\mu\nu} + \mathcal{L}_\mathrm{F} + \mathcal{L}_\mathrm{SQ}, \quad \mathcal{L}_\mathrm{SQ} = \mathrm{tr}\ (D_\mu \phi)^\dagger (D^\mu \phi) - m_\phi^2\ \mathrm{tr}\ \phi^\dagger \phi, \tag{1}$$

where $\mathcal{L}_\mathrm{F}$ denotes the quark part and m_ϕ the bare mass of scalar-quarks ϕ. In the actual calculation, we use a discretized Euclidean action on the $16^3 \times 32$ lattice at β=5.7,[9] i.e., lattice spacing a^{-1} $\simeq$1.1GeV.[11] The parameters employed in the analysis are summarized in Table 1.

The gauge-invariant local operators $O(\vec{x}, t)$ of scalar-quark hadrons and chimera hadrons are summarized in Table 2. We introduce "scalar-quark flavor" denoted by i, j, k and investigate the scalar-quark flavor non-singlet mesons, which do not have disconnected diagrams in their correlators. Note that, without the "scalar-quark flavor" degrees of freedom, the baryonic local operators of $\phi\phi\phi$ and $\phi\phi\psi$ inevitably

Table 2. Summary table of new-type hadrons. The indices i, j and k denote the scalar-quark flavor degrees of freedom. Γ_M^{ij}, Γ_B^{ijk} and Γ_B^{ij} are some tensors on the scalar-quark flavor.

Names		Lorentz properties	Local operators
Scalar-quark meson	$(\phi^\dagger\phi)$	Scalar	$M_s(x) = \Gamma_M^{ij} \phi_a^{\dagger i}(x) \phi_a^j(x)$
Scalar-quark baryon	$(\phi\phi\phi)$	Scalar	$B_s(x) = \Gamma_B^{ijk} \epsilon_{abc} \phi_a^i(x) \phi_b^j(x) \phi_c^k(x)$
Chimera meson	$(\phi^\dagger\psi)$	Spinor	$C_M^\alpha(x) = \phi_a^\dagger(x) \psi_a^\alpha(x)$
Chimera baryon	$(\psi\psi\phi)$	Scalar	$C_B(x) = \epsilon_{abc} ({\psi_a}^T(x) C\gamma_5 \psi_b(x)) \phi_c(x)$
Chimera baryon	$(\phi\phi\psi)$	Spinor	$C_B^\alpha(x) = \Gamma_B^{ij} \epsilon_{abc} \phi_a^i(x) \phi_b^j(x) \psi_c^\alpha(x)$



Fig. 1. (a) "Scalar-quark meson" mass squared $M^2_{\phi^\dagger\phi}$ and (b) "scalar-quark baryon" mass squared $M^2_{\phi\phi\phi}$ plotted against the bare scalar-quark mass squared m^2_ϕ at $a^{-1} \simeq 1.1$GeV.

vanish due to the anti-symmetric tensor ϵ_{abc}. We calculate the temporal correlator $G(t) \equiv \frac{1}{V}\sum_{\vec{x}}\langle O(\vec{x},t)O^\dagger(\vec{0},0)\rangle$, where the total momentum is projected to be zero. The mass M of these hadrons are obtained as $M \simeq -\frac{1}{T}\ln G(T)$ for large T.

Here, we show the lattice results for the masses of new-type hadrons. Figure 1 shows the squared scalar-quark-meson mass $M^2_{\phi^\dagger\phi}$ and the squared scalar-quark-baryon mass squared $M^2_{\phi\phi\phi}$, plotted against the bare scalar-quark mass squared m^2_ϕ at $a^{-1} \simeq 1.1$GeV. Even for zero bare scalar-quark mass, scalar-quark hadrons have a large mass as $M_{\phi^\dagger\phi} \simeq 3$GeV and $M_{\phi\phi\phi} \simeq 4.7$GeV. We find the "constituent scalar-quark picture", i.e., $M_{\phi^\dagger\phi} \simeq 2M_\phi$ and $M_{\phi\phi\phi} \simeq 3M_\phi$, where $M_\phi \simeq (1.5-1.6)$GeV is the constituent scalar-quark mass. The calculation can be performed even in the region $m^2_\phi < 0$ due to large quantum corrections on ϕ. We also find the relations, $M^2_{\phi^\dagger\phi} \simeq 4m^2_\phi + \text{const.}$ and $M^2_{\phi\phi\phi} \simeq 9m^2_\phi + \text{const.}$ from the figure. Together with the "constituent scalar-quark picture", we reach the relation $M^2_\phi \simeq m^2_\phi + \Sigma_\phi$, where Σ_ϕ is the self-energy of ϕ and is expected to be insensitive to m_ϕ. This is a natural relation between the renormalized mass and the bare mass for scalar particles.

Chimera hadrons also have a large mass even at $m_\phi = m_\psi = 0$, i.e., $M_{\phi^\dagger\psi} \simeq 1.9$GeV for chimera mesons $\phi^\dagger\psi$, and $M_{\psi\psi\phi} \simeq 2.2$GeV, $M_{\phi\phi\psi} \simeq 3.6$GeV for chimera baryons $(\psi\psi\phi, \phi\phi\psi)$. We find a "constituent scalar-quark/quark picture", i.e., an approximate relation as $M_{m\phi+n\psi} \simeq mM_\phi + nM_\psi$ with the constituent quark mass $M_\psi \simeq 0.4$GeV, and the large constituent scalar-quark mass $M_\phi \simeq (1.5-1.6)$GeV.[9]



(a)Chimera meson Φ†Ψ

(b)Chimera baryon ΦΦΨ

Fig. 2. The conjectured wave-function of the quark ψ around the heavy scalar-quarks in the chimera hadrons. Similar to chimera mesons, the wave-function of ψ in chimera baryons distributes around the two ϕ's localized near the center of mass.

From the m_ψ-dependence of chimera hadron masses, we conjecture a similar structure between chimera mesons $\phi^\dagger\psi$ and chimera baryons $\phi\phi\psi$.[9] The wave-function of ψ in a chimera meson $\phi^\dagger\psi$ is distributed around the heavy scalar-quark $\phi^\dagger$ due to the large mass of ϕ, and, similarly, the wave-function of ψ in a chimera baryon $\phi\phi\psi$ is distributed around the point-like "di-scalar-quark" $\phi\phi$. (See Fig. 2.)

3. Summary and Conclusion

We have performed the first study of light "scalar-quarks" ϕ (colored scalar particles or idealized diquarks) and their color-singlet hadronic states in quenched SU$(3)_c$ lattice QCD in terms of dynamical mass generation. We have investigated the mass of "scalar-quark mesons" $\phi^\dagger\phi$, "scalar-quark baryons" $\phi\phi\phi$ and "chimera hadrons" ($\phi^\dagger\psi$, $\psi\psi\phi$, $\phi\phi\psi$), which are composed of quarks ψ and scalar-quarks ϕ. We have observed the large dynamical mass generation of scalar-quarks ϕ about 1.5GeV at $a^{-1} \simeq 1.1$GeV due to large quantum corrections by gluons, even at the zero bare scalar-quark mass $m_\phi = 0$. This lattice result also indicates that plausible diquarks used in effective hadron models cannot be described as the point-like particles and should have a much larger size than $a \simeq 0.2$fm.[9,10]

This study indicates that, even without χSB, large dynamical mass generation in the strong interaction occurs for the scalar-quark systems. Together with the large glueball mass and the large difference between the current and the constituent charm-quark masses, this type of mass generation would generally occur in the strong interaction, and therefore we conjecture that all colored particles generally acquire a large effective mass due to dressed gluon effects,[9,10] as shown in Fig. 3.



Fig. 3. Schematic figure for dynamical mass generation of colored particles. Even without chiral symmetry breaking, colored particles generally acquire a large effective mass due to dressed gluons.

References

1. Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* **122**, 345 (1961); *ibid.* **124**, 246 (1961).
2. H. Iida, M. Oka and H. Suganuma, *Eur. Phys. J. A* **23**, 305 (2005).
3. J. E. Mandula and M. Ogilvie, *Phys. Lett. B* **185**, 127 (1987).
4. K. Amemiya and H. Suganuma, *Phys. Rev. D* **60**, 114509 (1999).
5. C. J. Morningstar and M. Peardon, *Phys. Rev. D* **60**, 034509 (1999).
6. N. Ishii, H. Suganuma and H. Matsufuru, *Phys. Rev. D* **66**, 094506 (2002).
7. Particle Data Group (W.M. Yao *et al.*), *J. Phys. G* **33**, 1 (2006).
8. A. De Rujula, H. Georgi and S. L. Glashow, *Phys. Rev. D* **12**, 147 (1975).
9. H. Iida, H. Suganuma and T. T. Takahashi, *Phys. Rev. D* **75**, 114503 (2007).
10. H. Iida, H. Suganuma and T. T. Takahashi, *AIP Conf. Proc.* **915**, 256 (2007).
11. T. T. Takahashi *et al.*, *Phys. Rev. Lett.* **86**, 18 (2001); *Phys. Rev. D* **65**, 114509 (2002).